

SEQIRUS FLU VACCINE QUOTATION REQUEST FORM FOR 2019/20 SEASON



To receive your FLUAD[®] quotation, please complete this form and return to the Seqirus team at fl.salesuk@seqirus.com or alternatively call 0345 0093 804.

Vaccine name	Licensed for (prescribing information overleaf)	NHS list price (ex. VAT)	Please provide number of doses required for 2019/20 season
FLUAD[®] (influenza vaccine, adjuvanted)	Adults aged 65 years and over	£9.79	<input type="text"/>

CONTACT INFORMATION (PLEASE USE CAPITAL LETTERS WHEN COMPLETING THIS FORM)

Account name <input type="text"/>	Contact name <input type="text"/>
Address <input type="text"/>	Best day/time to contact <input type="text"/>
Town <input type="text"/>	Other information <input type="text"/>
Postcode <input type="text"/>	
Telephone <input type="text"/>	
Email address <input type="text"/>	NHS practice code <input type="text"/>
	Date <input type="text"/>

A Seqirus sales representative will contact you shortly to discuss your vaccine offer and will be able to place your order.

FLUAD® PRESCRIBING INFORMATION

Fluad® suspension for injection in pre-filled syringe Influenza Vaccine (surface antigen, inactivated, adjuvanted with MF59C.1)

Presentation: Each 0.5ml of Fluad® contains 15 micrograms of each of three purified virus strains that comply with the World Health Organization trivalent vaccine recommendations (Northern Hemisphere) for the current season, with adjuvant MF59C.1 (9.75mg squalene, 1.175mg polysorbate 80, 1.175mg sorbitan trioleate, 0.66mg sodium citrate, 0.04mg citric acid, water). **Indications:** Active immunisation against influenza in the elderly (65 years of age and over), especially for those with an increased risk of associated complications. **Dosage and Administration:** Adults aged 65 years and over: single 0.5ml dose by intramuscular injection into the deltoid muscle using a 1-inch needle. **Contra-indications:** Hypersensitivity to the active substances, components of the adjuvant, excipients, to chicken or egg proteins (such as ovalbumin), kanamycin, neomycin sulphate, formaldehyde, cetyltrimethylammonium bromide (CTAB), barium sulphate, hydrocortisone or in anyone who has had an anaphylactoid reaction to previous influenza vaccination. Immunisation should be postponed in patients with febrile illness or acute infection. **Warnings and Precautions:** Appropriate medical treatment and supervision should be readily available in case of an anaphylactic event following administration. Do not inject intravascularly or subcutaneously. Endogenous or iatrogenic immunosuppression may result in insufficient antibody response. **Latex-sensitive individuals:** Although no natural rubber latex is detected in the syringe tip cap, the safe use of Fluad® in latex-sensitive individuals has not been established. **Interactions:** No clinical data on concomitant administration with other vaccines are available. If Fluad® needs to be used at the same time as another vaccine, immunisation should be carried out on separate limbs. It should be noted that the adverse reactions may be intensified. **Pregnancy and Lactation:** Not applicable. **Effects on ability to drive and use machines:** Fluad® has no or negligible

influence on the ability to drive and use machines. **Side Effects:** The most common reactions are headache, myalgia, injection site pain and tenderness, fatigue, nausea, diarrhoea, vomiting, sweating, arthralgia, fever, malaise, and shivering; local reactions include redness, swelling, ecchymosis, and induration. Uncommon reactions include rash. The following have been reported post-marketing: thrombocytopenia, lymphadenopathy, asthenia, influenza-like illness, extensive swelling of injected limb, injection-site cellulitis-like reaction, allergic reactions including anaphylactic shock (in rare cases), anaphylaxis, angioedema, vasculitis with transient renal involvement, and neurological disorders such as encephalomyelitis, Guillain-Barré syndrome, convulsions, neuritis, neuralgia, paraesthesia, syncope, and presyncope. **Overdose:** Overdosage is unlikely to have any untoward effect. **Legal Category:** POM. **Package Quantities:** Packs of 1 or 10 pre-filled syringes. **Marketing Authorisation Number:** UK: PL 46752/0001. **Basic NHS Cost:** £9.79 per 0.5ml pre-filled syringe, £97.90 per 10 pack. **Marketing Authorisation Holder:** Seqirus S.r.l., Via Fiorentina 1, 53100 Siena, Italy

For full prescribing information and details of other side effects see the Summary of Product Characteristics at www.medicines.org.uk/emc

Date of preparation: August 2018 UK/FLUD/0818/0098

© Registered trade mark Seqirus S.r.l, Italy

Adverse events should be reported. Reporting forms and information can be found at www.mhra.gov.uk/yellowcard. Adverse events relating to Seqirus products should also be reported to Seqirus UK Limited on 01748 828816